

4



4

Flavomycin®

ANTIBIOTIC
Type A
Medicated Article

For increased rate of weight gain and improved feed efficiency in broiler chickens, growing-finishing swine and growing turkeys.

ACTIVE DRUG INGREDIENTS

Bambermycins 4 grams per pound (8.8 grams per kilogram)

INGREDIENTS

Roughage products and/or calcium carbonate, dried bambermycins fermentation product, ethoxyquin (a preservative), mineral oil.

MIXING DIRECTIONS AND PERMITTED CLAIMS ON REVERSE SIDE

FOR USE IN MANUFACTURED FEEDS ONLY

Not For Human Use
Restricted Drug (California) -
Use only as directed

Net Weight – 50 pounds
(22.68 kg)

Flavomycin® (bambermycins)

Flavomycin®

(bambermycins)

ANTIBIOTIC Type A Medicated Article

PERMITTED CLAIMS AND LIMITATIONS

SPECIES	DRUG	LEVEL/TON	CLAIM(S)	LIMITATIONS
Broiler Chickens	Bambermycins	1-2 Grams*	Increased rate of weight gain; improved feed efficiency	None
Growing-Finishing Swine	Bambermycins	2 Grams	Increased rate of weight gain; improved feed efficiency	None
	Bambermycins	2-4 Grams*	Increased rate of weight gain	
Growing Turkeys	Bambermycins	2 Grams	Increased rate of weight gain; improved feed efficiency	None
	Bambermycins	1-2 Grams*	Increased feed efficiency	

*Only a single level of bambermycins (i.e., 4 grams/ton bambermycins) can be indicated on a feed tag. A different feed tag must be submitted for each level of bambermycins needed for your complete feeding program.

MIXING DIRECTIONS:

It is recommended that Flavomycin® 4 be diluted with a suitable grain carrier before addition to the final feed. A dilution of 1 part of Flavomycin® 4 and 9 parts grain carrier is the suggested working premix. The table below shows premix and working premix addition levels for various commonly used feeds:

FEED	BAMBERMYCINS LEVEL	POUNDS PREMIX/TON FLAVOMYCIN® 4	1-9 WORKING PREMIX (POUNDS/TON)
Broiler Chickens	1 gram/ton	.25	2.50
	2 grams/ton	.50	5.00
Growing-Finishing Swine	2 grams/ton	.50	5.00
	3 grams/ton	.75	7.50
	4 grams/ton	1.00	10.00
Growing Turkeys	1 gram/ton	.25	2.50
	2 grams/ton	.50	5.00

Thoroughly mix both working premix and finished feed to ensure complete and uniform distribution of the Flavomycin®.

FEEDING DIRECTIONS:

Feed containing Flavomycin® should be fed continuously as the sole ration.

Distributed by:
HUVEPHARMA, INC.
Peachtree City, GA 30269

 **HUVEPHARMA®**
NADA # 44-759, Approved by FDA

5040-02
Rev. 01-2013

236059 H-235002-A 2-13

FREIGHT SHIPPING BAG
MEETING REQUIREMENTS OF
APPLICABLE FREIGHT CLASSIFICATION
GUARANTEED BY
BANCROFT BAG, INC.
WEST MONROE, LOUISIANA

18x7.5x33